



Huntington's Disease Society of America

The information provided by speakers in workshops, forums, sharing/networking sessions and any other educational presentation made as part of the 2009 HDSA convention program is for informational use only.

HDSA encourages all attendees to consult with their primary care provider, neurologist or other healthcare provider about any advice, exercise, medication, treatment, nutritional supplement or regimen that may have been mentioned as part of any presentation.

Presenter Disclosures

Samuel Frank, MD

The following personal financial relationships with commercial interests relevant to this presentation existed during the past 12 months:

Consultant to Lundbeck
Speaker for Allergan




Huntington's Disease Society of America

Progression of Huntington's Disease

Samuel Frank, M.D.
Boston University School of Medicine
Assistant Professor of Neurology

Additional Disclosure

- This talk is not for the faint of heart



Overview

- The starting point
- The finish line
- The many years between
 - Reasons for decline
 - Maintaining independence
 - Improving quality of life

Gene Test – CAG Repeats

- Accurate gene test for individuals has been available since 1993
- 98.9% sensitivity
- Median number of repeats is 44
 - Range from 36-121
- Number of repeats does not determine course or severity of disease, but does generally correlates with age at onset.

Early Disease Features

- Chorea
- Personality changes
- Behavioral problems
- Clumsiness
- Subtle walking changes
- Mild speech rate and rhythm problems
- Memory problems
- In retrospect, often have preceding mood disorder, decline at work, or relationship difficulties.

Advanced Disease

- Stiffness and slowness of movement
 - More prominent than chorea, but chorea can also re-emerge
- Dystonia (Abnormal fixed postures)
- Weight loss
- Dementia
- Rate of progression varies, but duration of disease ranges from 10-25+ years

Causes of Death

- Causes of death are typically related to loss of movement:
 - Pneumonia
 - Aspiration
 - Nutrition
 - Chronic skin ulcers
- Suicide
 - ~ 25% attempt at least once
 - 8-9% of all deaths due to completed suicide
 - Risk factors include childlessness, depression, single, living alone, other suicides in family


HD Staging

- Based on 13 point Total Functional Capacity (TFC) score
 - Occupation (3 points)
 - Finances (3 points)
 - Domestic Chores (2 points)
 - ADL (3 points)
 - Care Level (2 points)
- Stage 1 = 11-13, 2 = 7-10, 3 = 3-6, 4 = 1-2, 5 = 0
- Typically lose ~1/2 point per year
- Not good for children/young adults, those with prominent psychiatric disease, or late stages.

Measuring Function

- TFC
- Functional checklist
 - 25 items
 - Ability or inability to do tasks
- Independence scale
 - 0-100% of 'normal' function
 - 5-10% increments

Changing Disease Progression



Disease-Modifying Strategies

- None proven to slow disease
- CoEnzyme Q10 (CARE-HD)
 - 13% slowing of the staging scale decline
 - 2CARE underway
- Creatine
- Others
 - Minocycline, blueberry extract, trehalose, omega 3 fatty acids (fish oil, ethyl-EPA), antidepressants (SSRI's)

What Is Associated with Disability?

- Early HD
 - Not studied well or extensively
- Late HD
 - Motor and cognitive difficulties

What is Associated with Disability?

- Although one of the most asked questions by patients, not well studied.
- Studies to date conclude:
 - Cognitive dysfunction is an important determinant of disability
 - Motor effects tend to be more disabling later
 - Treating motor aspects of HD tends to have little impact, but needs further study
- However, most studies examine mid-stage disease and not pre-symptomatic or early disease

I Shoulson, 1981

- 10 men, 12 women studied
- Treatment of chorea and depression out of proportion to physical disability
- In the first 7 years of HD, occupation capacities and financial management skills were most affected
- Improvement in chorea and depression was temporary and did not change functional outcome

R Mayeux, 1986

- Followed 33 patients over 'several years'
- Intellectual impairment and depression correlated with reduced functional capacity.
 - When somatic symptoms eliminated, no longer significantly associated
- There was no association of functional disability with:
 - Duration of illness
 - Motor disability
 - Age at onset

A Feigin, 1995

- Followed 129 people
- As functional capacity worsened, chorea lessened and dystonia intensified.
- No correlation between rate of functional decline and:
 - age at onset of HD
 - body weight
 - gender of affected parent
 - history of neuroleptic use.

K Marder, 2000

- 960 patients were followed for 18 months
- TFC declined by 0.72 units/year, faster for those with score of 7-13 at baseline
- Associated with less rapid decline in TFC:
 - Longer disease duration
 - Better cognitive status at baseline
- Associated with more rapid decline in Independent Scale:
 - Depression symptoms
- No effect:
 - Age at onset of HD, sex, weight, or education

N Mahant, 2003

- 1,026 patients followed for median of 2.7 years
 - From HSG UHDRS database
- All stages of disease studied
- Conclusions:
 - Rate of progression was associated with younger age at onset
 - ?CAG repeat length implications
 - Chorea was associated with weight loss, but chorea and dystonia were not major determinants of disability
 - Based on scale totals

JM Hamilton, 2003

- 22 patients
- Apathy and executive dysfunction behavioral index was strongly related to decline in ADL's
- Behavioral dysfunction may impede ability to use motor or cognitive skills that remain available in early HD
- JC Rothlind et al, 1993 also examined higher level of control and dysfunction in early HD:
 - Psychomotor speed
 - The ability to regulate attention

V Wheelock, 2003

- Used 3,070 patients with clinically definite HD in HSG UHDRS database
 - 228 resided in nursing homes
- For 1,559 patients, there was longitudinal data for 1.9 years
 - 87 patients moved from home to nursing home

V Wheelock, 2003: Findings

- NH residents had worse motor function:
 - Chorea, bradykinesia, gait abnormality, and imbalance
- More likely to have behavioral disease:
 - Obsessions, compulsions, delusions, auditory hallucinations
 - More aggressive, disruptive and irritable behaviors

V Wheelock, 2003

- Predictive of NH placement:
 - Impaired gait
 - Impaired tandem walking
 - Bradykinesia
- Important conclusion: motor variables alone predicted NH placement

S Frank, TBA

- CARE-HD database
- 347 participants at baseline
- Association examined between chorea, other motor items and TFC and Functional Checklist (totals and individual items)
- Disabilities associated with chorea:
 - Employment
 - Driving
 - Caring for children
 - Doing housework

S Frank, TBA

- TFC total was also associated with severity of chorea
- Individual items from TFC associated with chorea:
 - Occupation
 - Domestic chores

What Is Needed?

- COHORT
- The "Framingham Heart Study" of HD
- Prospectively follow:
 - Symptomatic HD
 - Gene positive, pre-symptomatic
 - At risk
 - Controls

Treatments - Symptomatic

- Movement disorders
- Psychiatric treatments
- Non-pharmacologic interventions
- All medications must be reconsidered with advancing disease

Treatments - Movements


- Chorea
 - Medications
 - Must weigh benefit of reducing chorea vs. side effects
- Slowness and Stiffness
 - Anti-Parkinsonian drugs
- Dystonia
 - Botulinum toxin
 - Medications

Treatments - Psychiatric

- Depression
 - Antidepressants
- Hallucinations, abnormal beliefs
 - Neuroleptics
 - May have benefit on movements also
 - May cause weight gain
- Sleep
 - Behavioral measures
 - Hypnotics
 - Disruption of the house is one of the main determinants of nursing home placement in Alzheimer's disease

Treatment Strategies: Cognitive Disorder

- **Memory**
 - Recognition easier than recall
 - Provide hints, yes/no questions
- **Learning**
 - Provide simple & concrete instructions
 - Limit distractions
- **Organization**
 - Help pts with lists, calendar, and routines
- **Bradyphrenia**
 - Do not interpret a lack of response as a "NO" response
 - Have patience



Conclusions

- In early HD, occupation, finances and driving may be affected
- Attention and cognitive changes may impact functioning early in disease
- Motor features (specifically chorea) may impact function early in HD
- A better understanding of HD may help with future treatments

Thank you!

